

A short HAP tutorial

**(A more comprehensive tutorial is
available here**

and

A related book is available here

and

The HAP home page is here)

Graham Ellis

Contents

1	Simplicial complexes & CW complexes	4
1.1	The Klein bottle as a simplicial complex	4
1.2	The Quillen complex	5
1.3	The Quillen complex as a reduced CW-complex	5
1.4	Constructing a regular CW-complex from its face lattice	6
1.5	Cup products	7
1.6	CW maps and induced homomorphisms	8
2	Cubical complexes & permutahedral complexes	10
2.1	Cubical complexes	10
2.2	Permutahedral complexes	11
2.3	Constructing pure cubical and permutahedral complexes	13
2.4	Computations in dynamical systems	13
3	Covering spaces	15
3.1	Cellular chains on the universal cover	15
3.2	Spun knots and the Satoh tube map	16
3.3	Cohomology with local coefficients	18
3.4	Distinguishing between two non-homeomorphic homotopy equivalent spaces	19
3.5	Second homotopy groups of spaces with finite fundamental group	19
3.6	Third homotopy groups of simply connected spaces	20
4	Topological data analysis	22
4.1	Persistent homology	22
4.2	Mapper clustering	23
4.3	Digital image analysis	24
5	Group theoretic computations	25
5.1	Third homotopy group of a suspension of an Eilenberg-MacLane space	25
5.2	Representations of knot quandles	25
5.3	Aspherical 2-complexes	26
5.4	Bogomolov multiplier	26
6	Cohomology of groups	27
6.1	Finite groups	27
6.2	Nilpotent groups	28
6.3	Crystallographic groups	29

6.4	Arithmetic groups	29
6.5	Artin groups	29
6.6	Graphs of groups	30
7	Cohomology operations	31
7.1	Steenrod operations on the classifying space of a finite 2-group	31
7.2	Steenrod operations on the classifying space of a finite p -group	33
8	Bredon homology	34
8.1	Davis complex	34
8.2	Arithmetic groups	34
8.3	Crystallographic groups	35
9	Simplicial groups	36
9.1	Crossed modules	36
9.2	Eilenberg-MacLane spaces	37
10	Parallel computation	38
10.1	An embarassingly parallel computation	38
	References	39

Chapter 1

Simplicial complexes & CW complexes

1.1 The Klein bottle as a simplicial complex

The following example constructs the Klein bottle as a simplicial complex K on 9 vertices, and then constructs the cellular chain complex $C_* = C_*(K)$ from which the integral homology groups $H_1(K, \mathbb{Z}) = \mathbb{Z}_2 \oplus \mathbb{Z}$, $H_2(K, \mathbb{Z}) = 0$ are computed. The chain complex $D_* = C_* \otimes_{\mathbb{Z}} \mathbb{Z}_2$ is also constructed and used to compute the mod-2 homology vector spaces $H_1(K, \mathbb{Z}_2) = \mathbb{Z}_2 \oplus \mathbb{Z}_2$, $H_2(K, \mathbb{Z}_2) = \mathbb{Z}_2$. Finally, a presentation $\pi_1(K) = \langle x, y : yxy^{-1}x \rangle$ is computed for the fundamental group of K .

Example

```
gap> 2simplices:=
> [[1,2,5], [2,5,8], [2,3,8], [3,8,9], [1,3,9], [1,4,9],
> [4,5,8], [4,6,8], [6,8,9], [6,7,9], [4,7,9], [4,5,7],
> [1,4,6], [1,2,6], [2,6,7], [2,3,7], [3,5,7], [1,3,5]];;
gap> K:=SimplicialComplex(2simplices);
Simplicial complex of dimension 2.

gap> C:=ChainComplex(K);
Chain complex of length 2 in characteristic 0 .

gap> Homology(C,1);
[ 2, 0 ]
gap> Homology(C,2);
[ ]

gap> D:=TensorWithIntegersModP(C,2);
Chain complex of length 2 in characteristic 2 .

gap> Homology(D,1);
2
gap> Homology(D,2);
1

gap> G:=FundamentalGroup(K);
<fp group of size infinity on the generators [ f1, f2 ]>
gap> RelatorsOfFpGroup(G);
[ f2*f1*f2~-1*f1 ]
```

1.2 The Quillen complex

Given a group G one can consider the partially ordered set $\mathcal{A}_p(G)$ of all non-trivial elementary abelian p -subgroups of G , the partial order being set inclusion. The order complex $\Delta\mathcal{A}_p(G)$ is a simplicial complex which is called the *Quillen complex*.

The following example constructs the Quillen complex $\Delta\mathcal{A}_2(S_7)$ for the symmetric group of degree 7 and $p = 2$. This simplicial complex involves 11291 simplices, of which 4410 are 2-simplices..

Example

```
gap> K:=QuillenComplex(SymmetricGroup(7),2);
Simplicial complex of dimension 2.

gap> Size(K);
11291

gap> K!.nrSimplices(2);
4410
```

1.3 The Quillen complex as a reduced CW-complex

Any simplicial complex K can be regarded as a regular CW complex. Different datatypes are used in HAP for these two notions. The following continuation of the above Quillen complex example constructs a regular CW complex Y isomorphic to (i.e. with the same face lattice as) $K = \Delta\mathcal{A}_2(S_7)$. An advantage to working in the category of CW complexes is that it may be possible to find a CW complex X homotopy equivalent to Y but with fewer cells than Y . The cellular chain complex $C_*(X)$ of such a CW complex X is computed by the following commands. From the number of free generators of $C_*(X)$, which correspond to the cells of X , we see that there is a single 0-cell and 160 2-cells. Thus the Quillen complex $\Delta\mathcal{A}_2(S_7) \simeq \bigvee_{i=1}^{160} S^2$ has the homotopy type of a wedge of 160 2-spheres. This homotopy equivalence is given in [Kso00, (15.1)] where it was obtained by purely theoretical methods.

Example

```
gap> Y:=RegularCWComplex(K);
Regular CW-complex of dimension 2

gap> C:=ChainComplex(Y);
Chain complex of length 2 in characteristic 0 .

gap> C!.dimension(0);
1
gap> C!.dimension(1);
0
gap> C!.dimension(2);
160
```

Note that for regular CW complexes Y the function `ChainComplex(Y)` returns the cellular chain complex $C_*(X)$ of a (typically non-regular) CW complex X homotopy equivalent to Y . The cellular chain complex $C_*(Y)$ of Y itself can be obtained as follows.

Example

```
gap> CC:=ChainComplexOfRegularCWComplex(Y);
Chain complex of length 2 in characteristic 0 .

gap> CC!.dimension(0);
1316
gap> CC!.dimension(1);
5565
gap> CC!.dimension(2);
4410
```

1.4 Constructing a regular CW-complex from its face lattice

The following example begins by creating a 2-dimensional annulus A as a regular CW-complex, and testing that it has the correct integral homology $H_0(A, \mathbb{Z}) = \mathbb{Z}$, $H_1(A, \mathbb{Z}) = \mathbb{Z}$, $H_2(A, \mathbb{Z}) = 0$.

Example

```
gap> FL:=[]; #The face lattice
gap> FL[1]:=[[1,0],[1,0],[1,0],[1,0]];
gap> FL[2]:=[[2,1,2],[2,3,4],[2,1,4],[2,2,3],[2,1,4],[2,2,3]];
gap> FL[3]:=[[4,1,2,3,4],[4,1,2,5,6]];
gap> FL[4]:=[];
gap> A:=RegularCWComplex(FL);
Regular CW-complex of dimension 2

gap> Homology(A,0);
[ 0 ]
gap> Homology(A,1);
[ 0 ]
gap> Homology(A,2);
[ ]
```

Next we construct the direct product $Y = A \times A \times A \times A \times A$ of five copies of the annulus. This is a 10-dimensional CW complex involving 248832 cells. It will be homotopy equivalent $Y \simeq X$ to a CW complex X involving fewer cells. The CW complex X may be non-regular. We compute the cochain complex $D_* = \text{Hom}_{\mathbb{Z}}(C_*(X), \mathbb{Z})$ from which the cohomology groups

$$\begin{aligned}
 H^0(Y, \mathbb{Z}) &= \mathbb{Z}, \\
 H^1(Y, \mathbb{Z}) &= \mathbb{Z}^5, \\
 H^2(Y, \mathbb{Z}) &= \mathbb{Z}^{10}, \\
 H^3(Y, \mathbb{Z}) &= \mathbb{Z}^{10}, \\
 H^4(Y, \mathbb{Z}) &= \mathbb{Z}^5, \\
 H^5(Y, \mathbb{Z}) &= \mathbb{Z}, \\
 H^6(Y, \mathbb{Z}) &= 0
 \end{aligned}$$

are obtained.

Example

```
gap> Y:=DirectProduct(A,A,A,A,A);
Regular CW-complex of dimension 10
```

```

gap> Size(Y);
248832
gap> C:=ChainComplex(Y);
Chain complex of length 10 in characteristic 0 .

gap> D:=HomToIntegers(C);
Cochain complex of length 10 in characteristic 0 .

gap> Cohomology(D,0);
[ 0 ]
gap> Cohomology(D,1);
[ 0, 0, 0, 0, 0 ]
gap> Cohomology(D,2);
[ 0, 0, 0, 0, 0, 0, 0, 0, 0, 0 ]
gap> Cohomology(D,3);
[ 0, 0, 0, 0, 0, 0, 0, 0, 0, 0 ]
gap> Cohomology(D,4);
[ 0, 0, 0, 0, 0 ]
gap> Cohomology(D,5);
[ 0 ]
gap> Cohomology(D,6);
[ ]

```

1.5 Cup products

Continuing with the previous example, we consider the first and fifth generators $g_1^1, g_5^1 \in H^1(W, \mathbb{Z}) = \mathbb{Z}^5$ and establish that their cup product $g_1^1 \cup g_5^1 = -g_7^2 \in H^2(W, \mathbb{Z}) = \mathbb{Z}^{10}$ is equal to minus the seventh generator of $H^2(W, \mathbb{Z})$. We also verify that $g_5^1 \cup g_1^1 = -g_1^1 \cup g_5^1$.

Example

```

gap> cup11:=CupProduct(FundamentalGroup(Y));
function( a, b ) ... end

gap> cup11([1,0,0,0,0],[0,0,0,0,1]);
[ 0, 0, 0, 0, 0, 0, -1, 0, 0, 0 ]

gap> cup11([0,0,0,0,1],[1,0,0,0,0]);
[ 0, 0, 0, 0, 0, 0, 1, 0, 0, 0 ]

```

This computation of low-dimensional cup products is achieved using group-theoretic methods to approximate the diagonal map $\Delta: Y \rightarrow Y \times Y$ in dimensions ≤ 2 . In order to construct cup products in higher degrees HAP requires a cellular inclusion $\bar{Y} \hookrightarrow Y \times Y$ with projection $p: \bar{Y} \rightarrow Y$ that induces isomorphisms on integral homology. The function `DiagonalApproximation(Y)` constructs a candidate inclusion, but the projection $p: \bar{Y} \rightarrow Y$ needs to be tested for homology equivalence. If the candidate inclusion passes this test then the function `CupProduct(Y)`, involving the candidate space, can be used for cup products.

The following example calculates $g_3^3 \cup g_3^1 = g_1^4$ where $W = S \times S \times S \times S$ is the direct product of four circles, and where g_k^n denotes the k -th generator of $H^n(W, \mathbb{Z})$.

Example

```

gap> S:=SimplicialComplex([[1,2],[2,3],[1,3]]);;
gap> S:=RegularCWComplex(S);;
gap> W:=DirectProduct(S,S,S,S);;
gap> cup:=CupProduct(W);
function( p, q, vv, ww ) ... end

gap> cup(3,1,[0,0,1,0],[0,0,1,0]);
[ 1 ]

#Now test that the diagonal construction is valid.
gap> D:=DiagonalApproximation(W);;
gap> p:=D!.projection;
Map of regular CW-complexes

gap> P:=ChainMap(p);
Chain Map between complexes of length 4 .

gap> IsIsomorphismOfAbelianFpGroups(Homology(P,0));
true
gap> IsIsomorphismOfAbelianFpGroups(Homology(P,1));
true
gap> IsIsomorphismOfAbelianFpGroups(Homology(P,2));
true
gap> IsIsomorphismOfAbelianFpGroups(Homology(P,3));
true
gap> IsIsomorphismOfAbelianFpGroups(Homology(P,4));
true

```

1.6 CW maps and induced homomorphisms

A *strictly cellular* map $f: X \rightarrow Y$ of regular CW-complexes is a cellular map for which the image of any cell is a cell (of possibly lower dimension). Inclusions of CW-subcomplexes, and projections from a direct product to a factor, are examples of such maps. Strictly cellular maps can be represented in HAP, and their induced homomorphisms on (co)homology and on fundamental groups can be computed.

The following example begins by visualizing the trefoil knot $\kappa \in \mathbb{R}^3$. It then constructs a regular CW structure on the complement $Y = D^3 \setminus \text{Nbhd}(\kappa)$ of a small tubular open neighbourhood of the knot lying inside a large closed ball D^3 . The boundary of this tubular neighbourhood is a 2-dimensional CW-complex B homeomorphic to a torus $\mathbb{S}^1 \times \mathbb{S}^1$ with fundamental group $\pi_1(B) = \langle a, b : aba^{-1}b^{-1} = 1 \rangle$. The inclusion map $f: B \hookrightarrow Y$ is constructed. Then a presentation $\pi_1(Y) = \langle x, y \mid xy^{-1}x^{-1}yx^{-1}y^{-1} \rangle$ and the induced homomorphism $\pi_1(B) \rightarrow \pi_1(Y)$, $a \mapsto y^{-1}xy^{-1}$, $b \mapsto y$ are computed. This induced homomorphism is an example of a *peripheral system* and is known to contain sufficient information to characterize the knot up to ambient isotopy.

Finally, it is verified that the induced homology homomorphism $H_2(B, \mathbb{Z}) \rightarrow H_2(Y, \mathbb{Z})$ is an isomorphism.

Example

```
gap> K:=PureCubicalKnot(3,1);;
gap> ViewPureCubicalKnot(K);;
```

Example

```
gap> K:=PureCubicalKnot(3,1);;
gap> f:=KnotComplementWithBoundary(ArcPresentation(K));
Map of regular CW-complexes

gap> G:=FundamentalGroup(Target(f));
<fp group of size infinity on the generators [ f1, f2 ]>
gap> RelatorsOfFpGroup(G);
[ f1*f2^-1*f1^-1*f2*f1^-1*f2^-1 ]

gap> F:=FundamentalGroup(f);
[ f1, f2 ] -> [ f2^-1*f1*f2^2*f1*f2^-1, f1 ]

gap> phi:=ChainMap(f);
Chain Map between complexes of length 2 .

gap> H:=Homology(phi,2);
[ g1 ] -> [ g1 ]
```

Chapter 2

Cubical complexes & permutahedral complexes

2.1 Cubical complexes

A *finite simplicial complex* can be defined to be a CW-subcomplex of the canonical regular CW-structure on a simplex Δ^n of some dimension n . Analogously, a *finite cubical complex* is a CW-subcomplex of the regular CW-structure on a cube $[0, 1]^n$ of some dimension n . Equivalently, but more conveniently, we can replace the unit interval $[0, 1]$ by an interval $[0, k]$ with CW-structure involving $2k + 1$ cells, namely one 0-cell for each integer $0 \leq j \leq k$ and one 1-cell for each open interval $(j, j + 1)$ for $0 \leq j \leq k - 1$. A *finite cubical complex* M is a CW-subcomplex $M \subset [0, k_1] \times [0, k_2] \times \cdots \times [0, k_n]$ of a direct product of intervals, the direct product having the usual direct product CW-structure. The equivalence of these two definitions follows from the Gray code embedding of a mesh into a hypercube. We say that the cubical complex has *ambient dimension* n . A cubical complex M of ambient dimension n is said to be *pure* if each cell lies in the boundary of an n -cell. In other words, M is pure if it is a union of unit n -cubes in \mathbb{R}^n , each unit cube having vertices with integer coordinates.

HAP has a datatype for finite cubical complexes, and a slightly different datatype for pure cubical complexes.

The following example constructs the granny knot (the sum of a trefoil knot with its reflection) as a 3-dimensional pure cubical complex, and then displays it.

Example

```
gap> K:=PureCubicalKnot(3,1);
prime knot 1 with 3 crossings

gap> L:=ReflectedCubicalKnot(K);
Reflected( prime knot 1 with 3 crossings )

gap> M:=KnotSum(K,L);
prime knot 1 with 3 crossings + Reflected( prime knot 1 with 3 crossings )

gap> Display(M);
```

Next we construct the complement $Y = D^3 \setminus \overset{\circ}{M}$ of the interior of the pure cubical complex M . Here D^3 is a rectangular region with $M \subset \overset{\circ}{D^3}$. This pure cubical complex Y is a union of 5891 unit

3-cubes. We contract Y to get a homotopy equivalent pure cubical complex YY consisting of the union of just 775 unit 3-cubes. Then we convert YY to a regular CW-complex W involving 11939 cells. We contract W to obtain a homotopy equivalent regular CW-complex WW involving 5993 cells. Finally we compute the fundamental group of the complement of the granny knot, and use the presentation of this group to establish that the Alexander polynomial $P(x)$ of the granny is

$$P(x) = x^4 - 2x^3 + 3x^2 - 2x + 1 .$$

Example

```

gap> Y:=PureComplexComplement(M);
Pure cubical complex of dimension 3.

gap> Size(Y);
5891

gap> YY:=ZigZagContractedComplex(Y);
Pure cubical complex of dimension 3.

gap> Size(YY);
775

gap> W:=RegularCWComplex(YY);
Regular CW-complex of dimension 3

gap> Size(W);
11939

gap> WW:=ContractedComplex(W);
Regular CW-complex of dimension 2

gap> Size(WW);
5993

gap> G:=FundamentalGroup(WW);
<fp group of size infinity on the generators [ f1, f2, f3 ]>

gap> AlexanderPolynomial(G);
x_1^4-2*x_1^3+3*x_1^2-2*x_1+1

```

2.2 Permutahedral complexes

A finite pure cubical complex is a union of finitely many cubes in a tessellation of \mathbb{R}^n by unit cubes. One can also tessellate \mathbb{R}^n by permutahedra, and we define a finite n -dimensional pure *permutahedral complex* to be a union of finitely many permutahedra from such a tessellation. There are two features of pure permutahedral complexes that are particularly useful in some situations:

- Pure permutahedral complexes are topological manifolds with boundary.
- The method used for finding a smaller pure cubical complex M' homotopy equivalent to a given pure cubical complex M retains the homeomorphism type, and not just the homotopy type, of the space M .

To illustrate these features the following example begins by reading in a protein backbone from the online [Protein Database](#), and storing it as a pure cubical complex K . The ends of the protein have been joined, and the homology $H_i(K, \mathbb{Z}) = \mathbb{Z}$, $i = 0, 1$ is seen to be that of a circle. We can thus regard the protein as a knot $K \subset \mathbb{R}^3$. The protein is visualized as a pure permutahedral complex.

Example

```
gap> file:=HapFile("data1V2X.pdb");;
gap> K:=ReadPDBfileAsPurePermutahedralComplex("file");
Pure permutahedral complex of dimension 3.

gap> Homology(K,0);
[ 0 ]
gap> Homology(K,1);
[ 0 ]

Display(K);
```

An alternative method for seeing that the pure permutahedral complex K has the homotopy type of a circle is to note that it is covered by open permutahedra (small open neighbourhoods of the closed 3-dimensional permutahedral tiles) and to form the nerve $N = \text{Nerve}(\mathcal{U})$ of this open covering \mathcal{U} . The nerve N has the same homotopy type as K . The following commands establish that N is a 1-dimensional simplicial complex and display N as a circular graph.

Example

```
gap> N:=Nerve(K);
Simplicial complex of dimension 1.

gap> Display(GraphOfSimplicialComplex(N));
```

The boundary of the pure permutahedral complex K is a 2-dimensional CW-complex B homeomorphic to a torus. We next use the advantageous features of pure permutahedral complexes to compute the homomorphism

$$\phi: \pi_1(B) \rightarrow \pi_1(\mathbb{R}^3 \setminus \mathring{K}), a \mapsto yx^{-3}y^2x^{-2}yxy^{-1}, b \mapsto yx^{-1}y^{-1}x^2y^{-1}$$

where

$$\pi_1(B) = \langle a, b : aba^{-1}b^{-1} = 1 \rangle,$$

$$\pi_1(\mathbb{R}^3 \setminus \mathring{K}) \cong \langle x, y : y^2x^{-2}yxy^{-1} = 1, yx^{-2}y^{-1}x(xy^{-1})^2 = 1 \rangle.$$

Example

```
gap> Y:=PureComplexComplement(K);
Pure permutahedral complex of dimension 3.
gap> Size(Y);
418922

gap> YY:=ZigZagContractedComplex(Y);
Pure permutahedral complex of dimension 3.
gap> Size(YY);
3438

gap> W:=RegularCWComplex(YY);
Regular CW-complex of dimension 3

gap> f:=BoundaryMap(W);
```

Map of regular CW-complexes

```
gap> CriticalCells(Source(f));
[ [ 2, 1 ], [ 2, 261 ], [ 1, 1043 ], [ 1, 1626 ], [ 0, 2892 ], [ 0, 24715 ] ]

gap> F:=FundamentalGroup(f,2892);
[ f1, f2 ] -> [ f2*f1^-3*f2^2*f1^-2*f2*f1*f2^-1, f2*f1^-1*f2^-1*f1^2*f2^-1 ]

gap> G:=Target(F);
<fp group on the generators [ f1, f2 ]>
gap> RelatorsOfFpGroup(G);
[ f2^2*f1^-2*f2*f1*f2^-1, f2*f1^-2*f2^-1*f1*(f1*f2^-1)^2 ]
```

2.3 Constructing pure cubical and permutahedral complexes

An n -dimensional pure cubical or permutahedral complex can be created from an n -dimensional array of 0s and 1s. The following example creates and displays two 3-dimensional complexes.

Example

```
gap> A:=[[ [0,0,0], [0,0,0], [0,0,0] ],
>        [ [1,1,1], [1,0,1], [1,1,1] ],
>        [ [0,0,0], [0,0,0], [0,0,0] ]];;
gap> M:=PureCubicalComplex(A);
Pure cubical complex of dimension 3.

gap> P:=PurePermutahedralComplex(A);
Pure permutahedral complex of dimension 3.

gap> Display(M);
gap> Display(P);
```

2.4 Computations in dynamical systems

Pure cubical complexes can be useful for rigorous interval arithmetic calculations in numerical analysis. They can also be useful for trying to estimate approximations of certain numerical quantities. To illustrate the latter we consider the *Henon map*

$$f: \mathbb{R}^2 \rightarrow \mathbb{R}^2, \begin{pmatrix} x \\ y \end{pmatrix} \mapsto \begin{pmatrix} y + 1 - ax^2 \\ bx \end{pmatrix}.$$

Starting with $(x_0, y_0) = (0, 0)$ and iterating $(x_{n+1}, y_{n+1}) = f(x_n, y_n)$ with the parameter values $a = 1.4$, $b = 0.3$ one obtains a sequence of points which is known to be dense in the so called *strange attractor* \mathcal{A} of the Henon map. The first 10 million points in this sequence are plotted in the following example, with arithmetic performed to 100 decimal places of accuracy. The sequence is stored as a 2-dimensional pure cubical complex where each 2-cell is square of side equal to $\varepsilon = 1/500$.

Example

```
gap> M:=HenonOrbit([0,0],14/10,3/10,10^-7,500,100);
Pure cubical complex of dimension 2.
```

```
gap> Size(M);  
10287  
  
gap> Display(M);
```

Repeating the computation but with squares of side $\varepsilon = 1/1000$

Example

```
gap> M:=HenonOrbit([0,0],14/10,3/10,10^-7,1000,100);  
  
gap> Size(M);  
24949
```

we obtain the heuristic estimate

$$\delta \simeq \frac{\log 24949 - \log 10287}{\log 2} = 1.277$$

for the box-counting dimension of the attractor \mathcal{A} .

Chapter 3

Covering spaces

Let Y denote a finite regular CW-complex. Let \tilde{Y} denote its universal covering space. The covering space inherits a regular CW-structure which can be computed and stored using the datatype of a $\pi_1 Y$ -equivariant CW-complex. The cellular chain complex $C_*\tilde{Y}$ of \tilde{Y} can be computed and stored as an equivariant chain complex. Given an admissible discrete vector field on Y , we can endow Y with a smaller non-regular CW-structure whose cells correspond to the critical cells in the vector field. This smaller CW-structure leads to a more efficient chain complex $C_*\tilde{Y}$ involving one free generator for each critical cell in the vector field.

3.1 Cellular chains on the universal cover

The following commands construct a 6-dimensional regular CW-complex $Y \simeq S^1 \times S^1 \times S^1$ homotopy equivalent to a product of three circles.

Example

```
gap> A:=[[1,1,1],[1,0,1],[1,1,1]];;
gap> S:=PureCubicalComplex(A);;
gap> T:=DirectProduct(S,S,S);;
gap> Y:=RegularCWComplex(T);;
Regular CW-complex of dimension 6

gap> Size(Y);
110592
```

The CW-somplex Y has 110592 cells. The next commands construct a free $\pi_1 Y$ -equivariant chain complex $C_*\tilde{Y}$ homotopy equivalent to the chain complex of the universal cover of Y . The chain complex $C_*\tilde{Y}$ has just 8 free generators.

Example

```
gap> Y:=ContractedComplex(Y);;
gap> CU:=ChainComplexOfUniversalCover(Y);;
gap> List([0..Dimension(Y)],n->CU!.dimension(n));
[ 1, 3, 3, 1 ]
```

The next commands construct a subgroup $H < \pi_1 Y$ of index 50 and the chain complex $C_*\tilde{Y} \otimes_{\mathbb{Z}H} \mathbb{Z}$ which is homotopy equivalent to the cellular chain complex $C_*\tilde{Y}_H$ of the 50-fold cover \tilde{Y}_H of Y corresponding to H .

Example

```

gap> L:=LowIndexSubgroupsFpGroup(CU!.group,50);
gap> H:=L[Length(L)-1];
gap> Index(CU!.group,H);
50
gap> D:=TensorWithIntegersOverSubgroup(CU,H);
Chain complex of length 3 in characteristic 0 .

gap> List([0..3],D!.dimension);
[ 50, 150, 150, 50 ]

```

General theory implies that the 50-fold covering space \tilde{Y}_H should again be homotopy equivalent to a product of three circles. In keeping with this, the following commands verify that \tilde{Y}_H has the same integral homology as $S^1 \times S^1 \times S^1$.

Example

```

gap> Homology(D,0);
[ 0 ]
gap> Homology(D,1);
[ 0, 0, 0 ]
gap> Homology(D,2);
[ 0, 0, 0 ]
gap> Homology(D,3);
[ 0 ]

```

3.2 Spun knots and the Satoh tube map

We'll construct two spaces Y, W with isomorphic fundamental groups and isomorphic integral homology, and use the integral homology of finite covering spaces to establish that the two spaces have distinct homotopy types.

By *spinning* a link $K \subset \mathbb{R}^3$ about a plane $P \subset \mathbb{R}^3$ with $P \cap K = \emptyset$, we obtain a collection $Sp(K) \subset \mathbb{R}^4$ of knotted tori. The following commands produce the two tori obtained by spinning the Hopf link K and show that the space $Y = \mathbb{R}^4 \setminus Sp(K) = Sp(\mathbb{R}^3 \setminus K)$ is connected with fundamental group $\pi_1 Y = \mathbb{Z} \times \mathbb{Z}$ and homology groups $H_0(Y) = \mathbb{Z}$, $H_1(Y) = \mathbb{Z}^2$, $H_2(Y) = \mathbb{Z}^4$, $H_3(Y, \mathbb{Z}) = \mathbb{Z}^2$. The space Y is only constructed up to homotopy, and for this reason is 3-dimensional.

Example

```

gap> Hopf:=PureCubicalLink("Hopf");
Pure cubical link.

gap> Y:=SpunAboutInitialHyperplane(PureComplexComplement(Hopf));
Regular CW-complex of dimension 3

gap> Homology(Y,0);
[ 0 ]
gap> Homology(Y,1);
[ 0, 0 ]
gap> Homology(Y,2);
[ 0, 0, 0, 0 ]
gap> Homology(Y,3);

```

```

[ 0, 0 ]
gap> Homology(Y,4);
[ ]
gap> GY:=FundamentalGroup(Y);;
gap> GeneratorsOfGroup(GY);
[ f2, f3 ]
gap> RelatorsOfFpGroup(GY);
[ f3^-1*f2^-1*f3*f2 ]

```

An alternative embedding of two tori $L \subset \mathbb{R}^4$ can be obtained by applying the 'tube map' of Shin Satoh to a welded Hopf link [Sat00]. The following commands construct the complement $W = \mathbb{R}^4 \setminus L$ of this alternative embedding and show that W has the same fundamental group and integral homology as Y above.

Example

```

gap> L:=HopfSatohSurface();
Pure cubical complex of dimension 4.

gap> W:=ContractedComplex(RegularCWComplex(PureComplexComplement(L)));
Regular CW-complex of dimension 3

gap> Homology(W,0);
[ 0 ]
gap> Homology(W,1);
[ 0, 0 ]
gap> Homology(W,2);
[ 0, 0, 0, 0 ]
gap> Homology(W,3);
[ 0, 0 ]
gap> Homology(W,4);
[ ]

gap> GW:=FundamentalGroup(W);;
gap> GeneratorsOfGroup(GW);
[ f1, f2 ]
gap> RelatorsOfFpGroup(GW);
[ f1^-1*f2^-1*f1*f2 ]

```

Despite having the same fundamental group and integral homology groups, the above two spaces Y and W were shown by Kauffman and Martins [KFM08] to be not homotopy equivalent. Their technique involves the fundamental crossed module derived from the first three dimensions of the universal cover of a space, and counts the representations of this fundamental crossed module into a given finite crossed module. This homotopy inequivalence is recovered by the following commands which involves the 5-fold covers of the spaces.

Example

```

gap> CY:=ChainComplexOfUniversalCover(Y);
Equivariant chain complex of dimension 3
gap> LY:=LowIndexSubgroups(CY!.group,5);;
gap> invY:=List(LY,g->Homology(TensorWithIntegersOverSubgroup(CY,g),2));;

```

```

gap> CW:=ChainComplexOfUniversalCover(W);
Equivariant chain complex of dimension 3
gap> LW:=LowIndexSubgroups(CW!.group,5);
gap> invW:=List(LW,g->Homology(TensorWithIntegersOverSubgroup(CW,g),2));

gap> SSortedList(invY)=SSortedList(invW);
false

```

3.3 Cohomology with local coefficients

The $\pi_1 Y$ -equivariant cellular chain complex $C_*\tilde{Y}$ of the universal cover \tilde{Y} of a regular CW-complex Y can be used to compute the homology $H_n(Y,A)$ and cohomology $H^n(Y,A)$ of Y with local coefficients in a $\mathbb{Z}\pi_1 Y$ -module A . To illustrate this we consider the space Y arising as the complement of the trefoil knot, with fundamental group $\pi_1 Y = \langle x, y : xyx = yxy \rangle$. We take $A = \mathbb{Z}$ to be the integers with non-trivial $\pi_1 Y$ -action given by $x.1 = -1, y.1 = -1$. We then compute

$$\begin{aligned} H_0(Y,A) &= \mathbb{Z}_2, \\ H_1(Y,A) &= \mathbb{Z}_3, \\ H_2(Y,A) &= \mathbb{Z}. \end{aligned}$$

Example

```

gap> K:=PureCubicalKnot(3,1);
gap> Y:=PureComplexComplement(K);
gap> Y:=ContractedComplex(Y);
gap> Y:=RegularCWComplex(Y);
gap> Y:=SimplifiedComplex(Y);
gap> C:=ChainComplexOfUniversalCover(Y);
gap> G:=C!.group;
gap> GeneratorsOfGroup(G);
[ f1, f2 ]
gap> RelatorsOfFpGroup(G);
[ f2^-1*f1^-1*f2^-1*f1*f2*f1, f1^-1*f2^-1*f1^-1*f2*f1*f2 ]
gap> hom:=GroupHomomorphismByImages(G,Group([[ -1 ]]),[G.1,G.2],[[ -1 ]],[[ -1 ]]);
gap> A:=function(x); return Determinant(Image(hom,x)); end;;
gap> D:=TensorWithTwistedIntegers(C,A); #Here the function A represents
gap> #the integers with twisted action of G.
Chain complex of length 3 in characteristic 0 .
gap> Homology(D,0);
[ 2 ]
gap> Homology(D,1);
[ 3 ]
gap> Homology(D,2);
[ 0 ]

```

3.4 Distinguishing between two non-homeomorphic homotopy equivalent spaces

The granny knot is the sum of the trefoil knot and its mirror image. The reef knot is the sum of two identical copies of the trefoil knot. The following commands show that the degree 1 homology homomorphisms

$$H_1(p^{-1}(B), \mathbb{Z}) \rightarrow H_1(\tilde{X}_H, \mathbb{Z})$$

distinguish between the homeomorphism types of the complements $X \subset \mathbb{R}^3$ of the granny knot and the reef knot, where $B \subset X$ is the knot boundary, and where $p: \tilde{X}_H \rightarrow X$ is the covering map corresponding to the finite index subgroup $H < \pi_1 X$. More precisely, $p^{-1}(B)$ is in general a union of path components

$$p^{-1}(B) = B_1 \cup B_2 \cup \dots \cup B_t.$$

The function `FirstHomologyCoveringCokernels(f, c)` inputs an integer c and the inclusion $f: B \hookrightarrow X$ of a knot boundary B into the knot complement X . The function returns the ordered list of the lists of abelian invariants of cokernels

$$\text{coker}(H_1(p^{-1}(B_i), \mathbb{Z}) \rightarrow H_1(\tilde{X}_H, \mathbb{Z}))$$

arising from subgroups $H < \pi_1 X$ of index c . To distinguish between the granny and reef knots we use index $c = 6$.

Example

```
gap> K:=PureCubicalKnot(3,1);;
gap> L:=ReflectedCubicalKnot(K);;
gap> granny:=KnotSum(K,L);;
gap> reef:=KnotSum(K,K);;
gap> fg:=KnotComplementWithBoundary(ArcPresentation(granny));;
gap> fr:=KnotComplementWithBoundary(ArcPresentation(reef));;
gap> a:=FirstHomologyCoveringCokernels(fg,6);;
gap> b:=FirstHomologyCoveringCokernels(fr,6);;
gap> a=b;
false
```

3.5 Second homotopy groups of spaces with finite fundamental group

If $p: \tilde{Y} \rightarrow Y$ is the universal covering map, then the fundamental group of \tilde{Y} is trivial and the Hurewicz homomorphism $\pi_2 \tilde{Y} \rightarrow H_2(\tilde{Y}, \mathbb{Z})$ from the second homotopy group of \tilde{Y} to the second integral homology of \tilde{Y} is an isomorphism. Furthermore, the map p induces an isomorphism $\pi_2 \tilde{Y} \rightarrow \pi_2 Y$. Thus $H_2(\tilde{Y}, \mathbb{Z})$ is isomorphic to the second homotopy group $\pi_2 Y$.

If the fundamental group of Y happens to be finite, then in principle we can calculate $H_2(\tilde{Y}, \mathbb{Z}) \cong \pi_2 Y$. We illustrate this computation for Y equal to the real projective plane. The above computation shows that Y has second homotopy group $\pi_2 Y \cong \mathbb{Z}$.

Example

```
gap> K:=[ [1,2,3], [1,3,4], [1,2,6], [1,5,6], [1,4,5],
>        [2,3,5], [2,4,5], [2,4,6], [3,4,6], [3,5,6] ];;
gap> K:=MaximalSimplicesToSimplicialComplex(K);
Simplicial complex of dimension 2.
gap> Y:=RegularCWComplex(K);
```

```

Regular CW-complex of dimension 2
gap> # Y is a regular CW-complex corresponding to the projective plane.

gap> U:=UniversalCover(Y);
Equivariant CW-complex of dimension 2

gap> G:=U!.group;;
gap> # G is the fundamental group of Y, which by the next command
gap> # is finite of order 2.
gap> Order(G);
2

gap> U:=EquivariantCWComplexToRegularCWComplex(U,Group(One(G)));
Regular CW-complex of dimension 2
gap> #U is the universal cover of Y

gap> Homology(U,0);
[ 0 ]
gap> Homology(U,1);
[ ]
gap> Homology(U,2);
[ 0 ]

```

3.6 Third homotopy groups of simply connected spaces

For any path connected space Y with universal cover \tilde{Y} there is an exact sequence

$$\rightarrow \pi_4 \tilde{Y} \rightarrow H_4(\tilde{Y}, \mathbb{Z}) \rightarrow H_4(K(\pi_2 \tilde{Y}, 2), \mathbb{Z}) \rightarrow \pi_3 \tilde{Y} \rightarrow H_3(\tilde{Y}, \mathbb{Z}) \rightarrow 0$$

due to J.H.C. Whitehead. Here $K(\pi_2(\tilde{Y}), 2)$ is an Eilenberg-MacLane space with second homotopy group equal to $\pi_2 \tilde{Y}$.

3.6.1 First example

Continuing with the above example where Y is the real projective plane, we see that $H_4(\tilde{Y}, \mathbb{Z}) = H_3(\tilde{Y}, \mathbb{Z}) = 0$ since \tilde{Y} is a 2-dimensional CW-space. The exact sequence implies $\pi_3 \tilde{Y} \cong H_4(K(\pi_2 \tilde{Y}, 2), \mathbb{Z})$. Furthermore, $\pi_3 \tilde{Y} = \pi_3 Y$. The following commands establish that $\pi_3 Y \cong \mathbb{Z}$.

Example

```

gap> A:=AbelianPcpGroup([0]);
Pcp-group with orders [ 0 ]

gap> K:=EilenbergMacLaneSimplicialGroup(A,2,5);;
gap> C:=ChainComplexOfSimplicialGroup(K);
Chain complex of length 5 in characteristic 0 .

gap> Homology(C,4);
[ 0 ]

```

3.6.2 Second example

The following commands construct a 4-dimensional simplicial complex Y with 9 vertices and 36 4-dimensional simplices, and establish that

$$\pi_1 Y = 0, \pi_2 Y = \mathbb{Z}, H_3(Y, \mathbb{Z}) = 0, H_4(Y, \mathbb{Z}) = \mathbb{Z}, H_4(K(\pi_2 Y, 2), \mathbb{Z}) = \mathbb{Z}.$$

Example

```
gap> Y:= [ [ 1, 2, 4, 5, 6 ], [ 1, 2, 4, 5, 9 ], [ 1, 2, 5, 6, 8 ],
>         [ 1, 2, 6, 4, 7 ], [ 2, 3, 4, 5, 8 ], [ 2, 3, 5, 6, 4 ],
>         [ 2, 3, 5, 6, 7 ], [ 2, 3, 6, 4, 9 ], [ 3, 1, 4, 5, 7 ],
>         [ 3, 1, 5, 6, 9 ], [ 3, 1, 6, 4, 5 ], [ 3, 1, 6, 4, 8 ],
>         [ 4, 5, 7, 8, 3 ], [ 4, 5, 7, 8, 9 ], [ 4, 5, 8, 9, 2 ],
>         [ 4, 5, 9, 7, 1 ], [ 5, 6, 7, 8, 2 ], [ 5, 6, 8, 9, 1 ],
>         [ 5, 6, 8, 9, 7 ], [ 5, 6, 9, 7, 3 ], [ 6, 4, 7, 8, 1 ],
>         [ 6, 4, 8, 9, 3 ], [ 6, 4, 9, 7, 2 ], [ 6, 4, 9, 7, 8 ],
>         [ 7, 8, 1, 2, 3 ], [ 7, 8, 1, 2, 6 ], [ 7, 8, 2, 3, 5 ],
>         [ 7, 8, 3, 1, 4 ], [ 8, 9, 1, 2, 5 ], [ 8, 9, 2, 3, 1 ],
>         [ 8, 9, 2, 3, 4 ], [ 8, 9, 3, 1, 6 ], [ 9, 7, 1, 2, 4 ],
>         [ 9, 7, 2, 3, 6 ], [ 9, 7, 3, 1, 2 ], [ 9, 7, 3, 1, 5 ] ];;

gap> Y:=MaximalSimplicesToSimplicialComplex(Y);
Simplicial complex of dimension 4.

gap> Y:=RegularCWComplex(Y);
Regular CW-complex of dimension 4

gap> Order(FundamentalGroup(Y));
1
gap> Homology(Y,2);
[ 0 ]
gap> Homology(Y,3);
[ ]
gap> Homology(Y,4);
[ 0 ]
```

Whitehead's sequence reduces to an exact sequence

$$\mathbb{Z} \rightarrow \mathbb{Z} \rightarrow \pi_3 Y \rightarrow 0$$

in which the first map is $H_4(Y, \mathbb{Z}) = \mathbb{Z} \rightarrow H_4(K(\pi_2 Y, 2), \mathbb{Z}) = \mathbb{Z}$. In order to determine $\pi_3 Y$ it remains compute this first map. This computation is currently not available in HAP.

[The simplicial complex in this second example is due to W. Kihnel and T. F. Banchoff and is of the homotopy type of the complex projective plane. So, assuming this extra knowledge, we have $\pi_3 Y = 0$.]

Chapter 4

Topological data analysis

4.1 Persistent homology

Pairwise distances between 74 points from some metric space have been recorded and stored in a 74×74 matrix D . The following commands load the matrix, construct a filtration of length 100 on the first two dimensions of the associated clique complex (also known as the *Rips Complex*), and display the resulting degree 0 persistent homology as a barcode. A single bar with label n denotes n bars with common starting point and common end point.

```
Example
gap> file:=HapFile("data253a.txt");
gap> Read(file);

gap> G:=SymmetricMatrixToFilteredGraph(D,100);
Filtered graph on 74 vertices.

gap> K:=FilteredRegularCWComplex(CliqueComplex(G,2));
Filtered regular CW-complex of dimension 2

gap> P:=PersistentBettiNumbers(K,0);
gap> BarCodeCompactDisplay(P);
```

The next commands display the resulting degree 1 persistent homology as a barcode.

```
Example
gap> P:=PersistentBettiNumbers(K,1);
gap> BarCodeCompactDisplay(P);
```

The following command displays the 1 skeleton of the simplicial complex arising as the 65-th term in the filtration on the clique complex.

```
Example
gap> Y:=FiltrationTerm(K,65);
Regular CW-complex of dimension 1

gap> Display(HomotopyGraph(Y));
```

These computations suggest that the dataset contains two persistent path components (or clusters), and that each path component is in some sense periodic. The final command displays one possible representation of the data as points on two circles.

4.1.1 Background to the data

Each point in the dataset was an image consisting of 732×761 pixels. This point was regarded as a vector in $\mathbb{R}^{732 \times 761}$ and the matrix D was constructed using the Euclidean metric. The images were the following:

4.2 Mapper clustering

The following example reads in a set S of vectors of rational numbers. It uses the Euclidean distance $d(u, v)$ between vectors. It fixes some vector $u_0 \in S$ and uses the associated function $f: D \rightarrow [0, b] \subset \mathbb{R}, v \mapsto d(u_0, v)$. In addition, it uses an open cover of the interval $[0, b]$ consisting of 100 uniformly distributed overlapping open subintervals of radius $r = 29$. It also uses a simple clustering algorithm implemented in the function `cluster`.

These ingredients are input into the Mapper clustering procedure to produce a simplicial complex M which is intended to be a representation of the data. The complex M is 1-dimensional and the final command uses GraphViz software to visualize the graph. The nodes of this simplicial complex are "buckets" containing data points. A data point may reside in several buckets. The number of points in the bucket determines the size of the node. Two nodes are connected by an edge when their end-point nodes contain common data points.

Example

```
gap> file:=HapFile("data134.txt");;
gap> Read(file);
gap> dx:=EuclideanApproximatedMetric;;
gap> dz:=EuclideanApproximatedMetric;;
gap> L:=List(S,x->Maximum(List(S,y->dx(x,y))));;
gap> n:=Position(L,Minimum(L));;
gap> f:=function(x); return [dx(S[n],x)]; end;;
gap> P:=30*[0..100];; P:=List(P, i->[i]);;
gap> r:=29;;
gap> epsilon:=75;;
gap> cluster:=function(S)
> local Y, P, C;
> if Length(S)=0 then return S; fi;
> Y:=VectorsToOneSkeleton(S,epsilon,dx);
> P:=PiZero(Y);
> C:=Classify([1..Length(S)],P[2]);
> return List(C,x->S{x});
> end;;
gap> M:=Mapper(S,dx,f,dz,P,r,cluster);
Simplicial complex of dimension 1.

gap> Display(GraphOfSimplicialComplex(M));
```

4.2.1 Background to the data

The datacloud S consists of the 400 points in the plane shown in the following picture.

4.3 Digital image analysis

The following example reads in a digital image as a filtered pure cubical complex. The filtration is obtained by thresholding at a sequence of uniformly spaced values on the greyscale range. The persistent homology of this filtered complex is calculated in degrees 0 and 1 and displayed as two barcodes.

Example

```
gap> file:=HapFile("image1.3.2.png");
gap> F:=ReadImageAsFilteredPureCubicalComplex(file,20);
Filtered pure cubical complex of dimension 2.
gap> P:=PersistentBettiNumbers(F,0);
gap> BarCodeCompactDisplay(P);
```

Example

```
gap> P:=PersistentBettiNumbers(F,1);
gap> BarCodeCompactDisplay(P);
```

The 20 persistent bars in the degree 0 barcode suggest that the image has 20 objects. The degree 1 barcode suggests that 14 (or possibly 17) of these objects have holes in them.

4.3.1 Background to the data

The following image was used in the example.

Chapter 5

Group theoretic computations

5.1 Third homotopy group of a suspension of an Eilenberg-MacLane space

The following example uses the nonabelian tensor square of groups to compute the third homotopy group

$$\pi_3(S(K(G,1))) = \mathbb{Z}^{30}$$

of the suspension of the Eilenberg-MacLane space $K(G,1)$ for G the free nilpotent group of class 2 on four generators.

Example

```
gap> F:=FreeGroup(4);;G:=NilpotentQuotient(F,2);;
gap> ThirdHomotopyGroupOfSuspensionB(G);
[ 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0,
  0, 0, 0, 0, 0, 0, 0, 0 ]
```

5.2 Representations of knot quandles

The following example constructs the finitely presented quandles associated to the granny knot and square knot, and then computes the number of quandle homomorphisms from these two finitely presented quandles to the 17-th quandle in HAP's library of connected quandles of order 24. The number of homomorphisms differs between the two cases. The computation therefore establishes that the complement in \mathbb{R}^3 of the granny knot is not homeomorphic to the complement of the square knot.

Example

```
gap> Q:=ConnectedQuandle(24,17,"import");;
gap> K:=PureCubicalKnot(3,1);;
gap> L:=ReflectedCubicalKnot(K);;
gap> square:=KnotSum(K,L);;
gap> granny:=KnotSum(K,K);;
gap> gcsquare:=GaussCodeOfPureCubicalKnot(square);;
gap> gcgranny:=GaussCodeOfPureCubicalKnot(granny);;
gap> Qsquare:=PresentationKnotQuandle(gcsquare);;
gap> Qgranny:=PresentationKnotQuandle(gcgranny);;
gap> NumberOfHomomorphisms(Qsquare,Q);
408
```

```
gap> NumberOfHomomorphisms(Qgranny,Q);
24
```

5.3 Aspherical 2-complexes

The following example uses Polymake's linear programming routines to establish that the 2-complex associated to the group presentation $\langle x, y, z : xyx = yxy, yzy = zyz, xzx = zxz \rangle$ is aspherical (that is, has contractible universal cover). The presentation is Tietze equivalent to the presentation used in the computer code, and the associated 2-complexes are thus homotopy equivalent.

Example

```
gap> F:=FreeGroup(6);;
gap> x:=F.1;;y:=F.2;;z:=F.3;;a:=F.4;;b:=F.5;;c:=F.6;;
gap> rels:=[a^-1*x*y, b^-1*y*z, c^-1*z*x, a*x*(y*a)^-1,
> b*y*(z*b)^-1, c*z*(x*c)^-1];;
gap> Print(IsAspherical(F,rels),"\n");
Presentation is aspherical.

true
```

5.4 Bogomolov multiplier

The Bogomolov multiplier of a group is an isoclinism invariant. Using this property, the following example shows that there are precisely three groups of order 243 with non-trivial Bogomolov multiplier. The groups in question are numbered 28, 29 and 30 in GAP's library of small groups of order 243.

Example

```
gap> L:=AllSmallGroups(3^5);;
gap> C:=IsoclinismClasses(L);;
gap> for c in C do
> if Length(BogomolovMultiplier(c[1]))>0 then
> Print(List(c,g->IdGroup(g)),"\n\n\n"); fi;
> od;
[ [ 243, 28 ], [ 243, 29 ], [ 243, 30 ] ]
```

Chapter 6

Cohomology of groups

6.1 Finite groups

The following example computes the fourth integral cohomology of the Mathieu group M_{24} .

$$H^4(M_{24}, \mathbb{Z}) = \mathbb{Z}_{12}$$

Example

```
gap> GroupCohomology(MathieuGroup(24),4);  
[ 4, 3 ]
```

The following example computes the third integral homology of the Weyl group $W = \text{Weyl}(E_8)$, a group of order 696729600.

$$H_3(\text{Weyl}(E_8), \mathbb{Z}) = \mathbb{Z}_2 \oplus \mathbb{Z}_2 \oplus \mathbb{Z}_{12}$$

Example

```
p> L:=SimpleLieAlgebra("E",8,Rationals);;  
gap> W:=WeylGroup(RootSystem(L));;  
gap> Order(W);  
696729600  
gap> GroupHomology(W,3);  
[ 2, 2, 4, 3 ]
```

The preceding calculation could be achieved more quickly by noting that $W = \text{Weyl}(E_8)$ is a Coxeter group, and by using the associated Coxeter polytope. The following example uses this approach to compute the fourth integral homology of W . It begins by displaying the Coxeter diagram of W , and then computes

$$H_4(\text{Weyl}(E_8), \mathbb{Z}) = \mathbb{Z}_2 \oplus \mathbb{Z}_2 \oplus \mathbb{Z}_2 \oplus \mathbb{Z}_2.$$

Example

```
gap> D:=[[1, [2,3]], [2, [3,3]], [3, [4,3]], [5,3]], [5, [6,3]], [6, [7,3]], [7, [8,3]]];;  
gap> CoxeterDiagramDisplay(D);
```

Example

```
gap> polytope:=CoxeterComplex_alt(D,5);;  
gap> R:=FreeGResolution(polytope,5);  
Resolution of length 5 in characteristic 0 for <matrix group with  
8 generators> .
```


Example

```
gap> F:=FreeGroup(4);; N:=NilpotentQuotient(F,2);;
gap> GroupHomology(N,4);
[ 3, 3, 3, 3, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0,
  0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0,
  0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0,
  0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0 ]
```

6.3 Crystallographic groups

The following example computes

$$H_5(G, \mathbb{Z}) = \mathbb{Z}_2 \oplus \mathbb{Z}_2$$

for the 3-dimensional crystallographic space group G with Hermann-Mauguin symbol "P62"

Example

```
gap> GroupHomology(SpaceGroupBBNWZ("P62"),5);
[ 2, 2 ]
```

6.4 Arithmetic groups

The following example computes

$$H_6(SL_2(\mathcal{O}, \mathbb{Z}) = \mathbb{Z}_2$$

for \mathcal{O} the ring of integers of the number field $\mathbb{Q}(\sqrt{-2})$.

Example

```
gap> C:=ContractibleGcomplex("SL(2,0-2)");;
gap> R:=FreeGResolution(C,7);;
gap> Homology(TensorWithIntegers(R),6);
[ 2, 12 ]
```

6.5 Artin groups

The following example computes

$$H_5(G, \mathbb{Z}) = \mathbb{Z}_3$$

for G the classical braid group on eight strings.

Example

```
gap> D:=[[1, [2,3]], [2, [3,3]], [3, [4,3]], [4, [5,3]], [5, [6,3]], [6, [7,3]]];;
gap> CoxeterDiagramDisplay(D);;
```

Example

```
gap> R:=ResolutionArtinGroup(D,6);;
gap> C:=TensorWithIntegers(R);;
gap> Homology(C,5);
[ 3 ]
```

6.6 Graphs of groups

The following example computes

$$H_5(G, \mathbb{Z}) = \mathbb{Z}_2 \oplus \mathbb{Z}_2 \oplus \mathbb{Z}_2 \oplus \mathbb{Z}_2 \oplus \mathbb{Z}_2$$

for G the graph of groups corresponding to the amalgamated product $G = S_5 *_{S_3} S_4$ of the symmetric groups S_5 and S_4 over the canonical subgroup S_3 .

Example

```
gap> S5:=SymmetricGroup(5);SetName(S5,"S5");
gap> S4:=SymmetricGroup(4);SetName(S4,"S4");
gap> A:=SymmetricGroup(3);SetName(A,"S3");
gap> AS5:=GroupHomomorphismByFunction(A,S5,x->x);
gap> AS4:=GroupHomomorphismByFunction(A,S4,x->x);
gap> D:=[S5,S4,[AS5,AS4]];
gap> GraphOfGroupsDisplay(D);
```

Example

```
gap> R:=ResolutionGraphOfGroups(D,6);;
gap> Homology(TensorWithIntegers(R),5);
[ 2, 2, 2, 2, 2 ]
```

Chapter 7

Cohomology operations

7.1 Steenrod operations on the classifying space of a finite 2-group

The following example determines a presentation for the cohomology ring $H^*(Syl_2(M_{12}), \mathbb{Z}_2)$. The Lyndon-Hochschild-Serre spectral sequence, and Groebner basis routines from SINGULAR, are used to determine how much of a resolution to compute for the presentation.

Example

```
gap> G:=SylowSubgroup(MathieuGroup(12),2);;
gap> Mod2CohomologyRingPresentation(G);
Graded algebra GF(2)[ x_1, x_2, x_3, x_4, x_5, x_6, x_7 ] /
[ x_2*x_3, x_1*x_2, x_2*x_4, x_3^3+x_3*x_5,
  x_1^2*x_4+x_1*x_3*x_4+x_3^2*x_4+x_3^2*x_5+x_1*x_6+x_4^2+x_4*x_5,
  x_1^2*x_3^2+x_1*x_3*x_5+x_3^2*x_5+x_3*x_6,
  x_1^3*x_3+x_3^2*x_4+x_3^2*x_5+x_1*x_6+x_3*x_6+x_4*x_5,
  x_1*x_3^2*x_4+x_1*x_3*x_6+x_1*x_4*x_5+x_3*x_4^2+x_3*x_4*x_5+x_3*x_5^2\
2+x_4*x_6, x_1^2*x_3*x_5+x_1*x_3^2*x_5+x_3^2*x_6+x_3*x_5^2,
  x_3^2*x_4^2+x_3^2*x_5^2+x_1*x_5*x_6+x_3*x_4*x_6+x_4*x_5^2,
  x_1*x_3*x_4^2+x_1*x_3*x_4*x_5+x_1*x_3*x_5^2+x_3^2*x_5^2+x_1*x_4*x_6+\
x_2^2*x_7+x_2*x_5*x_6+x_3*x_4*x_6+x_3*x_5*x_6+x_4^2*x_5+x_4*x_5^2+x_6^2\
2, x_1*x_3^2*x_6+x_3^2*x_4*x_5+x_1*x_5*x_6+x_4*x_5^2,
  x_1^2*x_3*x_6+x_1*x_5*x_6+x_2^2*x_7+x_2*x_5*x_6+x_3*x_5*x_6+x_6^2
] with indeterminate degrees [ 1, 1, 1, 2, 2, 3, 4 ]
```

The command `CohomologicalData(G,n)` prints complete information for the cohomology ring $H^*(G, \mathbb{Z}_2)$ of a 2-group G provided that the integer n is at least the maximal degree of a relator in a minimal set of relators for the ring. Groebner basis routines from SINGULAR are called involved in the example.

The following example produces complete information on the Steenrod algebra of group number 8 in GAP's library of groups of order 32.

Example

```
Group number: 8
Group description: C2 . ((C4 x C2) : C2) = (C2 x C2) . (C4 x C2)

Cohomology generators
Degree 1: a, b
Degree 2: c, d
```

Degree 3: e
 Degree 5: f, g
 Degree 6: h
 Degree 8: p

Cohomology relations

1: f^2
 2: $c*h+e*f$
 3: $c*f$
 4: $b*h+c*g$
 5: $b*e+c*d$
 6: $a*h$
 7: $a*g$
 8: $a*f+b*f$
 9: $a*e+c^2$
 10: $a*c$
 11: $a*b$
 12: a^2
 13: $d*e*h+e^2*g+f*h$
 14: $d^2*h+d*e*f+d*e*g+f*g$
 15: $c^2*d+b*f$
 16: $b*c*g+e*f$
 17: $b*c*d+c*e$
 18: $b^2*g+d*f$
 19: b^2*c+c^2
 20: b^3+a*d
 21: $c*d^2*e+c*d*g+d^2*f+e*h$
 22: $c*d^3+d*e^2+d*h+e*f+e*g$
 23: $b^2*d^2+c*d^2+b*f+e^2$
 24: b^3*d
 25: $d^3*e^2+d^2*e*f+c^2*p+h^2$
 26: $d^4*e+b*c*p+e^2*g+g*h$
 27: $d^5+b*d^2*g+b^2*p+f*g+g^2$

Poincare series

$$(x^5+x^2+1)/(x^8-2*x^7+2*x^6-2*x^5+2*x^4-2*x^3+2*x^2-2*x+1)$$

Steenrod squares

$Sq^1(c)=0$
 $Sq^1(d)=b*b*b+d*b$
 $Sq^1(e)=c*b*b$
 $Sq^2(e)=e*d+f$
 $Sq^1(f)=c*d*b*b+d*d*b*b$
 $Sq^2(f)=g*b*b$
 $Sq^4(f)=p*a$
 $Sq^1(g)=d*d*d+g*b$
 $Sq^2(g)=0$
 $Sq^4(g)=c*d*d*d*b+g*d*b*b+g*d*d+p*a+p*b$
 $Sq^1(h)=c*d*d*b+e*d*d$
 $Sq^2(h)=d*d*d*b*b+c*d*d*d+g*c*b$
 $Sq^4(h)=d*d*d*d*b*b+g*e*d+p*c$
 $Sq^1(p)=c*d*d*d*b$

```
Sq^2(p)=d*d*d*d*b*b+c*d*d*d*d
Sq^4(p)=d*d*d*d*d*b*b+d*d*d*d*d*d+g*d*d*d*b+g*g*d+p*d*d
```

7.2 Steenrod operations on the classifying space of a finite p -group

The following example constructs the first eight degrees of the mod-3 cohomology ring $H^*(G, \mathbb{Z}_3)$ for the group G number 4 in GAP's library of groups of order 81. It determines a minimal set of ring generators lying in degree ≤ 8 and it evaluates the Bockstein operator on these generators. Steenrod powers for $p \geq 3$ are not implemented as no efficient method of implementation is known.

Example

```
gap> G:=SmallGroup(81,4);;
gap> A:=ModPSteenrodAlgebra(G,8);;
gap> List(ModPPringGenerators(A),x->Bockstein(A,x));
[ 0*v.1, 0*v.1, v.5, 0*v.1, (Z(3))*v.7+v.8+(Z(3))*v.9 ]
```

Chapter 8

Bredon homology

8.1 Davis complex

The following example computes the Bredon homology

$$\underline{H}_0(W, \mathcal{R}) = \mathbb{Z}^{21}$$

for the infinite Coxeter group W associated to the Dynkin diagram shown in the computation, with coefficients in the complex representation ring.

Example

```
gap> D:=[[1, [2, 3]], [2, [3, 3]], [3, [4, 3]], [4, [5, 6]]];;
gap> CoxeterDiagramDisplay(D);
```

Example

```
gap> C:=DavisComplex(D);;
gap> D:=TensorWithComplexRepresentationRing(C);;
gap> Homology(D,0);
[ 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0 ]
```

8.2 Arithmetic groups

The following example computes the Bredon homology

$$\underline{H}_0(SL_2(\mathcal{O}_{-3}), \mathcal{R}) = \mathbb{Z}_2 \oplus \mathbb{Z}^9$$

$$\underline{H}_1(SL_2(\mathcal{O}_{-3}), \mathcal{R}) = \mathbb{Z}$$

for \mathcal{O}_{-3} the ring of integers of the number field $\mathbb{Q}(\sqrt{-3})$, and \mathcal{R} the complex reflection ring.

Example

```
gap> R:=ContractibleGcomplex("SL(2,0-3)");;
gap> IsRigid(R);
false
gap> S:=BaryCentricSubdivision(R);;
gap> IsRigid(S);
true
gap> C:=TensorWithComplexRepresentationRing(S);;
gap> Homology(C,0);
[ 2, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0 ]
gap> Homology(C,1);
```

```
[ 0 ]
```

8.3 Crystallographic groups

The following example computes the Bredon homology

$$\underline{H}_0(G, \mathcal{R}) = \mathbb{Z}^{17}$$

for G the second crystallographic group of dimension 4 in GAP's library of crystallographic groups, and for \mathcal{R} the Burnside ring.

Example

```
gap> G:=SpaceGroup(4,2);;
gap> gens:=GeneratorsOfGroup(G);;
gap> B:=CrystGFullBasis(G);;
gap> R:=CrystGcomplex(gens,B,1);;
gap> IsRigid(R);
false
gap> S:=CrystGcomplex(gens,B,0);;
gap> IsRigid(S);
true
gap> D:=TensorWithBurnsideRing(S);;
gap> Homology(D,0);
[ 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0 ]
```

Chapter 9

Simplicial groups

9.1 Crossed modules

The following example concerns the crossed module

$$\partial: G \rightarrow \text{Aut}(G), g \mapsto (x \mapsto gxg^{-1})$$

associated to the dihedral group G of order 16. This crossed module represents, up to homotopy type, a connected space X with $\pi_i X = 0$ for $i \geq 3$, $\pi_2 X = Z(G)$, $\pi_1 X = \text{Aut}(G)/\text{Inn}(G)$. The space X can be represented, up to homotopy, by a simplicial group. That simplicial group is used in the example to compute

$$H_1(X, \mathbb{Z}) = \mathbb{Z}_2 \oplus \mathbb{Z}_2,$$

$$H_2(X, \mathbb{Z}) = \mathbb{Z}_2,$$

$$H_3(X, \mathbb{Z}) = \mathbb{Z}_2 \oplus \mathbb{Z}_2 \oplus \mathbb{Z}_2,$$

$$H_4(X, \mathbb{Z}) = \mathbb{Z}_2 \oplus \mathbb{Z}_2 \oplus \mathbb{Z}_2,$$

$$H_5(X, \mathbb{Z}) = \mathbb{Z}_2 \oplus \mathbb{Z}_2 \oplus \mathbb{Z}_2 \oplus \mathbb{Z}_2 \oplus \mathbb{Z}_2 \oplus \mathbb{Z}_2.$$

The simplicial group is obtained by viewing the crossed module as a crossed complex and using a nonabelian version of the Dold-Kan theorem.

Example

```
gap> C:=AutomorphismGroupAsCatOneGroup(DihedralGroup(16));
Cat-1-group with underlying group Group(
[ f1, f2, f3, f4, f5, f6, f7, f8, f9 ] ) .

gap> Size(C);
512
gap> Q:=QuasiIsomorph(C);
Cat-1-group with underlying group Group( [ f9, f8, f1, f2*f3, f5 ] ) .

gap> Size(Q);
32

gap> N:=NerveOfCatOneGroup(Q,6);
Simplicial group of length 6

gap> K:=ChainComplexOfSimplicialGroup(N);
Chain complex of length 6 in characteristic 0 .

gap> Homology(K,1);
[ 2, 2 ]
```

```

gap> Homology(K,2);
[ 2 ]
gap> Homology(K,3);
[ 2, 2, 2 ]
gap> Homology(K,4);
[ 2, 2, 2 ]
gap> Homology(K,5);
[ 2, 2, 2, 2, 2 ]

```

9.2 Eilenberg-MacLane spaces

The following example concerns the Eilenberg-MacLane space $X = K(\mathbb{Z}, 3)$ which is a path-connected space with $\pi_3 X = \mathbb{Z}$, $\pi_i X = 0$ for $3 \neq i \geq 1$. This space is represented by a simplicial group, and perturbation techniques are used to compute

$$H_7(X, \mathbb{Z}) = \mathbb{Z}_3.$$

Example

```

gap> A:=AbelianPcpGroup([0]);;AbelianInvariants(A);
[ 0 ]
gap> K:=EilenbergMacLaneSimplicialGroup(A,3,8);
Simplicial group of length 8

gap> C:=ChainComplexOfSimplicialGroup(K);
Chain complex of length 8 in characteristic 0 .

gap> Homology(C,7);
[ 3 ]

```

Chapter 10

Parallel computation

10.1 An embarrassingly parallel computation

The following example creates five child processes and uses them simultaneously to compute the second integral homology of each of the 267 groups of order 64. The final command shows that

$$H_2(G, \mathbb{Z}) = \mathbb{Z}_2^{15}$$

for the 267-th group G in GAP's library of small groups.

Example

```
gap> Processes:=List([1..5],i->ChildProcess());
gap> fn:=function(i);return GroupHomology(SmallGroup(64,i),2);end;;
gap> for p in Processes do
> ChildPut(fn,"fn",p);
> od;

gap> NrSmallGroups(64);
267

gap> L:=ParallelList([1..267],"fn",Processes);;

gap> L[267];
[ 2, 2, 2, 2, 2, 2, 2, 2, 2, 2, 2, 2, 2, 2, 2 ]
```

The function `ParallelList()` is built from HAP's six core functions for parallel computation.

References

- [KFM08] L. H. Kauffman and J. Faria Martins. Invariants of welded virtual knots via crossed module invariants of knotted surfaces. *Compos. Math.*, 144(4):1046–1080, 2008. [17](#)
- [Kso00] R. Ksontini. *Proprietes homotopiques du complexe de Quillen du groupe symetrique*. These de doctorat, Universitet de Lausanne, 2000. [5](#)
- [Sat00] S. Satoh. Virtual knot presentation of ribbon torus-knots. *J. Knot Theory Ramifications*, 9(4):531–542, 2000. [17](#)